

DIRECTIONS FOR USING BLAZE LIQUID WICK FUELS

STANDARD OPERATING INSTRUCTIONS

BLAZE PRODUCTS:

Our liquid wick products contain diethylene glycol (DEG). This product is ideal for warming foods, sauces, and beverages it should be used in chafing dishes only.

NEVER LIGHT THE FUEL WHILE HOLDING THE CAN.

NEVER LEAVE LIT BLAZE PRODUCTS UNATTENDED.

ALWAYS USE CAUTION WHEN HANDLING BLAZE PRODUCTS.

ALWAYS USE FLAME RETARDANT MATERIALS AROUND FUEL AREA

i.e. TABLECLOTH, NAPKINS AND TABLE SKIRTING.

DO NOT ALLOW TABLECLOTH AND TABLE DECORATIONS TO COME CLOSE TO THE PROXIMITY OF THE FLAME.

CAUTION:

BLAZE liquid wick products are combustible. It produces a HOT open flame.

Do not open or remove liquid from the container. May be harmful if swallowed. Avoid prolonged breathing of liquid vapor and over-exposure to skin and eyes.

STORAGE:

Store away from heat, sparks, or open flame

DIRECTIONS FOR USE:

Remove the water pan and food pan.

Place one or two cans of fuel under the chafing dish in the designated holder or tray. NEVER use more than *two* cans of fuel at a time under a chafing dish.

Wick is set for optimum heat. To remove cap, twist and pull. Save the cap for recapping.

Always begin with HOT water. Fill chafing dish water pan with hot water to indicated level marked on water pan. Check water pan periodically to ensure a sufficient water level is maintained.

NEVER USE WITHOUT WATER!

Light the product with a long stemmed match or long handled lighter.

TO EXTINGUISH:

Use metal snuffing device. **DO NOT BLOW OUT!** Let wick cool.

TIP:

If you have trouble lighting wick, simply invert can to allow wicking action to begin. Then light as usual. Do not throw away the can until all the fuel has been used. Recycle container.

FIRST AID:

If on skin thoroughly wash exposed area with soap and water. Remove contaminated clothing. Launder contaminated clothing before re-use. If in eyes flush with large amounts of water, lifting upper and lower lids occasionally. **GET MEDICAL ATTENTION.**

If swallowed and victim is conscious, give two glasses of water, induce vomiting immediately by sticking finger down throat. Call a physician. Never give anything by mouth to an unconscious person.

If inhaled move individual to fresh air.